



Solvipurity

ANALYTICAL LABORATORY · REYKJAVÍK, IS

SVP-2026-00344

ISSUED 2026-03-16 · ACCREDITATION AL-1142  
ISO/IEC 17025 · GMP · GLP

## CERTIFICATE OF ANALYSIS



AUTHENTIC

# Pal-AHK 200mg

Björn Healthcare ehf. · Sterile lyophilizate, 200 mg per 3 ml clear glass vial, rubber stopper + aluminium flip-off (white cake)

REPRESENTATIVE CHROMATOGRAM · HPLC-UV 205 NM



## BATCH NO.

BJRN-208STJV

## ANALYTICAL METHODS

RP-HPLC-UV 220 nm · LC-ESI-MS · AAA (amino acid analysis) · Ion chromatography (counter-ion) · Karl Fischer 2.5.32 · GC-MS (headspace) · ICP-MS · Kinetic chromogenic LAL 2.6.14 · Ph. Eur. 2.6.1 sterility · Ph. Eur. 2.6.12 microbial limits

## MANUFACTURED

2026-09-04

## EXPIRY

2028-06-04

## RECEIVED

2026-03-16

## RELEASE

2026-03-16

## DECLARED COMPOSITION

Pal-AHK 200mg — synthetic peptide; sequence: Palmitoyl-Ala-His-Lys; CAS n/a; theoretical MW 608.85 Da

## Analytical results

19 TESTS · ALL METHODS VALIDATED

| SUBSTANCE / PARAMETER                                       | RESULT   | LOQ | LIMIT                                   | METHOD                  |
|---|--|-----|---|-------------------------|
| ● Appearance — sterile lyophilized cake, white to off-white | Conforms   | —   | homogeneous white cake, no particulates | Visual                  |
| ● Solubility (water for injection, 2 mg/ml, 25 °C)          | Complete within 60 s — clear colourless solution | —   | Clear, no visible particles             | Visual (Ph. Eur. 2.2.1) |

| Identification — HPLC retention time               | Matches reference                | —           | ±2.0 % of ref                    | RP-HPLC-UV<br>220 nm<br>SVP-2026-00344 +        |
|--|----------------------------------|-------------|----------------------------------|---|
| ● Identification — sequence / mass match           | <b>Confirmed</b><br>CAS n/a      | —           | Match theoretical within ±1 Da   | LC-ESI-MS                                       |
| ● Molecular weight (measured)                      | <b>609.12 Da</b><br>Δ = +0.27 Da | 0.5 Da      | 608.85 Da ± 1.0 Da (theoretical) | ESI-MS  |
| ● Chromatographic purity (main peak)               | <b>98.53 %</b>                   | 0.05 %      | ≥ 98.0 %                         | RP-HPLC-UV<br>220 nm                            |
| ● Any single impurity (max)                        | <b>0.24 %</b>                    | 0.05 %      | ≤ 1.00 %                         | RP-HPLC-UV<br>220 nm                            |
| ● Peptide content (amino acid analysis)            | <b>85.8 % w/w</b>                | 0.5 %       | ≥ 80.0 % w/w                     | AAA (6 N HCl,<br>110 °C, 24 h)                  |
| ● Trifluoroacetate (TFA counter-ion)               | <b>0.32 % w/w</b>                | 0.05 %      | ≤ 1.00 % w/w                     | IC (ion chromatography)                         |
| ● Water content (Karl Fischer)                     | <b>3.26 % w/w</b>                | 0.1 %       | ≤ 5.0 % w/w                      | Ph. Eur.<br>2.5.32                              |
| ● Residual acetonitrile                            | <b>189 ppm</b>                   | 10 ppm      | ≤ 410 ppm (ICH Q3C Class 2)      | GC-MS<br>(headspace)                            |
| ● Residual DMF                                     | <b>117 ppm</b>                   | 10 ppm      | ≤ 880 ppm (ICH Q3C Class 2)      | GC-MS<br>(headspace)                            |
| ● Lead (Pb)  | <b>0.109 ppm</b>                 | 0.02 ppm    | ≤ 0.5 ppm (ICH Q3D parenteral)   | ICP-MS  |
| ● Arsenic + Cadmium + Mercury (total)              | <b>0.116 ppm</b>                 | 0.02 ppm    | ≤ 1.5 ppm (ICH Q3D parenteral)   | ICP-MS  |
| ● Bacterial endotoxins (LAL)                       | <b>4.52 EU/mg</b>                | 0.125 EU/mg | < 10.0 EU/mg                     | Kinetic chromogenic LAL (Ph. Eur. 2.6.14)       |
| ● TAMC (aerobic bacteria, pre-lyophilization bulk) | <b>1 CFU/g</b>                   | 1 CFU/g     | ≤ 10 <sup>2</sup> CFU/g          | Ph. Eur.<br>2.6.12                              |
| ● TYMC (yeast / molds, pre-lyophilization bulk)    | <b>5 CFU/g</b>                   | 1 CFU/g     | ≤ 10 <sup>1</sup> CFU/g          | Ph. Eur.<br>2.6.12                              |
| ● Sterility (final lyophilized vial)               | <b>Complies – no growth</b>      | —           | No growth, 14 d incubation       | Ph. Eur.<br>2.6.1 (direct inoculation)          |
| ● Container closure integrity                      | <b>Pass</b>                      | —           | No dye uptake                    | Dye ingress (0.05 % methylene blue, 2 h vacuum) |



